

DEPARTMENT OF WATER AND SANITATION

NO. 655

10 MAY 2019

**NATIONAL WATER ACT, 1998
(ACT NO.36 OF 1998)****PROPOSED CLASSES OF WATER RESOURCE AND RESOURCE QUALITY OBJECTIVES
FOR THE BERG CATCHMENT**

I, Gugile Nkwinti, in my capacity as Minister of Water and Sanitation and duly authorised in terms of Section 13(4) of the National Water Act, 1998 (Act No. 36 of 1998) hereby publish, the notice for the proposed classes of water resources and the proposed resource quality objectives for the Berg Catchment.

Any person who wishes to submit written comments with regard to the proposed classes of water resources and the proposed resource quality objectives should submit the comments within 60 days from the date of publication of this Notice to:

Director: Water Resource Classification
Attention: Ms Lebogang Matlala
Department of Water and Sanitation
Ndinaye Building 5046
178 Francis Baard Street
Private Bag x 313
Pretoria
0001
Facsimile: 012 336 6712
Email: matlalal@dws.gov.za



MR NKWINTI GE (MP)
MINISTER OF WATER AND SANITATION
DATE: 22/02/2019

SCHEDULE**DESCRIPTION OF THE WATER RESOURCE**

The proposed water resource classes and resource quality objectives are determined for all or part of every significant water resource as set out below:

Water Management Area:	Berg-Olifants Water Management Area
Drainage Region:	G1, G2 Secondary Drainage Region and G40A Quaternary Drainage Region
River(s):	The Berg River is the largest river in the study area, which also includes a number of smaller catchments within the City of Cape Town Metropolitan area such as the Diep, Kuils, Eerste, Lourens, Sir Lowry's, Steenbras, as well as various small catchments on the Cape Peninsula and along the West Coast.

A. PROPOSED WATER RESOURCE CLASSES AS REQUIRED IN TERMS OF SECTION 13(4)(a)(i)(aa) OF THE NATIONAL WATER ACT, 1998

- i. The proposed water resource classes for the Berg Catchment are listed in Table 1 according to the overall class per integrated unit of analysis (IUA), indicated in Figure 1.
- ii. IUAs are classified as either Class I: indicating high environmental protection and minimal utilisation; Class II indicating moderate protection and moderate utilisation; and Class III indicating sustainable minimal protection and high utilisation.
- iii. Table 1 provides the IUA, the recommended water resource class and its respective catchment configuration. The catchment configuration consists of a number of biophysical nodes representing river reaches or river resource units (RUs). The target ecological category (TEC) to be achieved or maintained for each RU in the IUA is provided.
- iv. It is important to note that additional existing geographically defined areas of specific ecological importance for water resources such as protected areas (e.g. Table Mountain National Park), critical biodiversity areas (CBAs), national freshwater environmental protection areas (NFEPA's) and the strategic water source areas (SWSA) should also be considered in terms of the recommended resource classes as these would indicate areas of specific importance that should be managed in a higher resource class (e.g. Class I) than would be the case for the average of all resource units across the IUA (e.g. in a Class II).

B. RESOURCE QUALITY OBJECTIVES OF WATER RESOURCES AS REQUIRED IN TERMS OF SECTION 13(4)(a)(i)(bb) OF THE NATIONAL WATER ACT, 1998

- i. Resource Quality Objectives (RQOs) are defined for prioritised RUs for each IUA in terms of water quantity, habitat and biota, and water quality. Prioritised RUs are indicated in Figure 1.
- ii. Table 2 to Table 10 provide the RQOs for RIVERS in priority RUs.
- iii. Table 11 to Table 17 provide the RQOs for ESTUARIES in priority RUs.
- iv. Table 18 provides the RQOs for DAMS in priority RUs.
- v. Table 19 provides the RQOs for GROUNDWATER in priority RUs.
- vi. RQOs will apply from the date signed off as determined in terms of Section 13(1) of the National Water Act, 1998, unless otherwise specified by the Minister.




Figure 1: Proposed Water Resource Classes for the Berg Catchment




Table 1: Summary of recommended Water Resource Classes for each IUA and the Target Ecological Category (TEC) for priority biophysical river and estuary nodes

Integrated Unit of Analysis (IUA)	Water Resource Class for IUA	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	% nMAR*
A1 Berg Estuary	II	G10M	A1-E01	Berg (Groot)	Bxi1	C	52
A2 Langebaan	II	G10M	A2-E04	Langebaan	Bxi3	A	N/A
D8 Upper Berg	II	G10A	D8-R01	Berg	Bvii13	A	98
		G10A	D8-R02	Berg	Bviii1	C	27
		G10C	D8-R03	Berg	Blii3	D	53
		G10C	D9-R04	Pombers	Bviii11	C	366
D9 Middle Berg	III	G10D	D9-R05	Kromme	Bvii3	D	89
		G10D	D9-R06	Berg	Bvii5	D	49
		G10E	C5-R07	Klein Berg	Blii4	C	82
		G10G	C5-R08	Vier-en-Twintig	Bi1	B/C	23
B4 Lower Berg	III	G10J	B4-R09	Berg	Bvii6	D	52
		G10K	B4-R10	Berg	Bvii12	D	51
		G21D	D10-R11	Diep	Bv1	D	66
		G21D	D10-R12	Diep	Biv6	D	68
D10 Diep	III	G21F	D10-E03	Rietvlei/ Diep	Bxi7	C	78
E11 Peninsula	II	G22B	E11-R13	Hout Bay	Bviii6	D	97
		G22A	E11-R14	Silvermine	Bvii20	C	98
		G22A	E11-E04	Wildevöelvlei	Bxi14	C	107
		G22D	E12-R15	Keyzers	Bvii7	D	93
E12 Cape Flats	III	G22K	E12-E05	Zandvlei	Bxi9	C	93
		G22K	E12-E05	Zeekoevlei	Bxi9	D	N/A
		G22F	D6-R16	Eerste (Jonkershoek)	Blii6	C	93
		G22G	D6-R17	Klippies	Biv8	D	77
D6 Eerste	III	G22H	D6-E06	Eerste	Bxi3	D	90
		G22J	D7-R18	Lourens	Bvii21	D	114
		G22K	D7-R19	Sir Lowry's Pass*	Bviii9	C	84
		G40A	D7-R20	Steenbras	Bvii22	B/C	81
D7 Sir Lowry's	II	G22J	D7-E07	Lourens	Bxi4	D	85

*Note: This is based on the estimated/simulated flow requirement in the system to meet downstream TECs as well as with current demands. Note that this will differ from the minimum flow requirement to meet the EWR at any given node. In some cases, the flow is above 100% of natural due to the impact of releases to meet downstream demands.

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
D8 Upper Berg	II	G10A	D8-R02	Berg River	Bviii1	C	Quantity	Low flows High flows	Indigenous riparian woody species	Upper zone cover abundance	Cover 25-60%																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
									Non-woody indigenous species		Cover 25-50%																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
									Reeds		No reeds																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
									Exotic species		Cover < 10%.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
									Terrestrial woody species		Cover < / = 15%.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
									Indigenous riparian woody species		Cover 25-50%																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
									Non-woody indigenous species		Cover 40-70%.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
								Biota	Fish	FRAI score	Fish condition	> 80% = B category																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																						
										Number of indigenous fish species.	Three species present: <i>Sandelia capensis</i> , <i>Galaxia zebratus</i> and <i>Pseudobarbus burgi</i>																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
										<i>Sandelia capensis</i>	FROC = 5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
										<i>Galaxias zebratus</i>	FROC = 5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
										<i>Pseudobarbus burgi</i>	FROC = 5																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
										Exotic fish species	No increase in the number of exotic fish present: <i>Onchorhynchus mykiss</i> (FROC = 5)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
										Invertebrates	Macroinvertebrate condition	MIRAI score	> 78 % = B/C category																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																					
							SASS5 and ASPT score	SASS5 score >180, ASPT ≥ 7.2.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
							Number of families	Diversity of invertebrate community	> / = 23 families, at an abundance of A to C.																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
							Quality	Salts	Nutrients			Maintenance low flows	Flows sufficient to maintain the river in a C category	Maintenance flows (million cubic metres)	High	0.000	2.143	Nov	1.071	Dec	0.803	Jan	0.726	Feb	0.803	Mar	1.296	Apr	2.679	May	4.147	Jun	4.285	Jul	4.285	Aug	3.888	Sep																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
												Maintenance high flows		Low	0.000	0.544	1.293	0.803	0.000	0.726	0.803	0.000	0.778	0.000	4.666	10.109	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
									Dissolved oxygen	for the maintenance of ecosystem health.	DO ≥ 8 milligrams per litre (5th percentile)
								Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Ideal category for full contact recreation.	≤ 130 counts/100ml (95th percentile)
								Geomorphology	D50	Sand particle size	0.521 > D50 > 0.319
									VEGRAI level 3 score.	Vegetation condition	> 62% = C category
									Exotic species		No exotic plant species.
									Terrestrial woody species		No terrestrial woody species.
									Indigenous riparian woody species	Marginal zone cover abundance	Cover < 10%.
									Non-woody indigenous species		Cover 50-75%.
							Habitat	Riparian vegetation	Reeds		No reeds
									Exotic species		Cover < 5%.
									Terrestrial woody species		Cover < 10%.
									Indigenous riparian woody species	Lower zone cover abundance	Cover 50-75%.
									Non-woody indigenous species		Cover 25-50%.
									Reeds		No reeds
									FRAI score	Fish condition	> 62% = C category
								Fish	Number of indigenous fish species.		One species present: <i>Sandelia capensis</i>
									<i>Sandelia capensis</i>	Indigenous species richness	FROC = 5
									Exotic fish species		No increase in the number of exotic fish present: <i>Micropterus dolomieu</i> (FROC = 5)
									MIRAI score	Macroinvertebrate condition	> 62%= C category
								Invertebrates	SASS5 and ASPT score	SASS scores	SASS5 score >134, ASPT ≥ 6.1.
									Number of families	Diversity of invertebrate community	> /= 21 families, at an abundance of A to C.

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																						
D8 Upper Berg	II	G10C	D8-R03	Berg River	Biii3	D	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a D category	Months	Low	0.003	5.803	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep							
											Maintenance flows (million cubic metres)	High	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
												≤ 0.075 milligrams/litre (50th percentile)	≤ 1.75 milligrams/litre (50th percentile)	≤ 55 milliSiemens/metre (95th percentile)	6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles)	2°C difference from ambient water temperature	DO ≥ 6 milligrams per litre (5th percentile)	≤ 0.073 milligrams per litre (95th percentile)	≤ 0.079 milligrams per litre (95th percentile)	≤ 0.0013 milligrams per litre (95th percentile)	≤ 4000 counts/100ml (95th percentile)	> 38% = D/E category	> 58% C/D category										
							Quality	System variables	Salts	Electrical conductivity (EC)	pH range	Water temperature	Dissolved oxygen	Ammonia	Atrazine	Endosulfan	Toxicity levels must not pose a threat to aquatic ecosystems.	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation.	Sediment particle size	Vegetation condition	Fish condition												
																						Pathogens	Escherichia coli	D16, D50, D84	VEGRAI level 3 score.	FRAI score							
																											Habitat	Geomorphology	Riparian vegetation	Fish			
							Biota	Fish	FRAI score	Fish condition																							

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric			
D9 Middle Berg	III	G10D	D9-R06	Berg River	Bvii5	D	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a D category	Maintenance flows (million cubic metres) High Low	Months	Sep	20.262
												Aug	25.299	
												Jul	37.63	
												Jun	24.388	
								May	0.000					
								Apr	10.152					
								Mar	2.585					
							Feb	2.342						
							Jan	2.621						
							Dec	2.648						
							Nov	5.200						
							Oct	14.246						
							0.000							
							≤ 0.125 milligrams/litre (50th percentile)							
							Quality	Nutrients	Phosphate (PO ₄ -P)	Nutrient levels must be maintained in the river at a eutrophic or better condition.	≤ 3.00 milligrams/litre (50th percentile)			
									Total inorganic nitrogen (TIN)					
									Electrical conductivity (EC)	Salt concentrations need to be maintained at present state levels.	95%tile ≤ 55 milliSiemens/metre EC			
								Salts	pH range	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles)			
									Water temperature		2°C difference from ambient water temperature			
									Dissolved oxygen		≥ 6 milligrams litre (5th percentile)			
									Toxins	Ammonia	Toxicity levels must not pose a threat to aquatic ecosystems.	≤ 0.073 milligrams per litre (95th percentile)		
Atrazine		≤ 0.079 milligrams per litre (95th percentile)												
			≤ 0.0013 milligrams per litre (95th percentile)											
Habitat	Geomorphology	GAI score -	Geomorphological condition	> 38% = D/E category										
		Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	> 18% = F category									
		Fish	FRAI score	Fish condition	> 22% = E category									
		Invertebrates	MIRAI score	Macroinvertebrate condition	> 78% = B/C category									
	Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation.	≤ 2500 counts/100ml (95th percentile)										
Toxins	Ammonia	Toxicity levels must not pose a threat to aquatic ecosystems.	≤ 0.073 milligrams per litre (95th percentile)											
	Atrazine		≤ 0.079 milligrams per litre (95th percentile)											
	Endusulfan		≤ 0.0013 milligrams per litre (95th percentile)											
System variables	Water temperature	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles)											
	Dissolved oxygen		2°C difference from ambient water temperature											
			DO ≥ 8 milligrams per litre (5th percentile)											
Salts	Electrical conductivity (EC)	Salt concentrations must be maintained in an Ideal category.	≤ 30 milliSiemens/metre (95th percentile)											

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
									Endosulfan		≤ 0.0013 milligrams per litre (95th percentile)
								Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation.	95 th tile ≤ 2500 cfu/100ml Escherichia coli
								Geomorphology	D50	Sand particle size	0.714 > D50 > 0.251
									VEGRAI level 3 score.	Vegetation condition	> 52% = D category
									Exotic species		No exotic plant species.
									Terrestrial woody species		No terrestrial woody species.
									Indigenous riparian woody species	Marginal zone cover abundance	Cover 50-75%.
									Non-woody indigenous species		Cover 15-25%.
									Reeds		No reeds
									Exotic species		Cover < 5%.
									Terrestrial woody species		Cover < 10%.
									Indigenous riparian woody species	Lower zone cover abundance	Cover 50-75%.
									Non-woody indigenous species		Cover 15-25%.
									Reeds		No reeds
									Exotic species		Cover < 10%.
									Terrestrial woody species		Cover < /= 15%.
									Indigenous riparian woody species	Upper zone cover abundance	Cover 50-75%.
									Non-woody indigenous species		Cover 10-20%
								Fish	FRAI score	Fish condition	> 52% = D category
									Exotic fish species	Indigenous species richness	No increase in the number of exotic fish present: <i>Cyprinus carpio</i> (FROC = 5), <i>Tilapia sparrmanii</i> , <i>Clarias gariepinus</i> , <i>Gambusia affinis</i>
								Invertebrates	MIRAI score	Macroinvertebrate condition	> 62% = C category
									SASS5 and ASPT score	SASS scores	SASS5 score >90, ASPT ≥ 4.6.
									Number of families	Diversity of invertebrate community	> /= 18 families, at an abundance of A to C.

Table 4: Resource Quality Objectives for RIVERS in priority Resource Units in the Integrated Unit of Analysis C5 Berg Tributaries

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																																		
C5 Berg Tributaries	II	G10F	C5-R07	Klein Berg River	Biii4	C	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a C category	Maintenance flows (million cubic metres)	Months	Low	1.422	1.110	Nov	0.754	Dec	0.000	Jan	0.398	Feb	0.305	Mar	0.291	Apr	0.338	May	1.002	Jun	1.391	Jul	1.744	Aug	1.619										
													High	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			
							Quality	Nutrients	Phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)	Nutrient levels must be maintained in the river at a mesotrophic or better condition.	≤ 0.075 milligrams/litre (50th percentile) ≤ 1.75 milligrams/litre (50th percentile)	Months	High	Oct	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000				
														High	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
								Salts	Electrical conductivity (EC)	Salt concentrations need to be maintained at levels that do not adversely affect aquatic ecosystems	≤ 55 milliSiemens/metre (95th percentile)	Months	Low	Oct	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000				
														High	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
								System variables	pH range Water temperature Dissolved oxygen	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles) 2°C difference from ambient water temperature ≥ 6 milligrams litre (5th percentile)	Months	Low	Oct	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			
														High	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
								Toxins	Ammonia Atrazine Endosulfan	Toxicity levels must not pose a threat to aquatic ecosystems.	≤ 0.073 milligrams per litre (95th percentile) ≤ 0.079 milligrams per litre (95th percentile) ≤ 0.0013 milligrams per litre (95th percentile)	Months	Low	Oct	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			
														High	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
							Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation.	≤ 2500 counts/100ml (95th percentile)	Months	Low	Oct	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			
													High	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
Habitat	Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	> 62% = C category	Months	Low	Oct	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000									
							High	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000							
Biota	Fish	FRAI score	Fish condition	> 58% = C/D category	Months	Low	Oct	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000									
							High	0.638	0.141	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000							
C5 Berg Tributaries	II	G10G	C5-R08	Vier-en-Twintig	Bi1	B/C	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a B/C category	Maintenance flows (million cubic metres)	Months	Low	2.050	1.631	Nov	1.115	Dec	0.000	Jan	0.731	Feb	0.563	Mar	0.573	Apr	0.674	May	1.811	Jun	2.510	Jul	3.886	Aug	0.748	Sep	1.497								
													High	0.646	0.217	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
							Habitat	Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	> 62% = C category	Months	Low	Oct	0.646	0.217	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			
														High	0.646	0.217	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
							Biota	Fish	FRAI score	Fish condition	> 58% = C/D category	Months	Low	Oct	0.646	0.217	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000			
														High	0.646	0.217	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
							Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a B/C category	Maintenance flows (million cubic metres)	Months	Low	Oct	0.646	0.217	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
														High	0.646	0.217	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
														High	0.646	0.217	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
														High	0.646	0.217	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Nutrients	Phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)	Nutrient levels must be maintained in the river at an oligotrophic condition.	≤ 0.025 milligrams per litre PO ₄ -P ≤ 0.70 milligrams per litre TIN
								Salts	Electrical conductivity (EC)	Salt concentrations need to be maintained in an Ideal category for aquatic ecosystems	≤ 30 milliSiemens/metre (95th percentile)
									pH range Water temperature Dissolved oxygen	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	4.5 ≤ pH ≤ 7.0 (5th and 95th percentiles) 2°C difference from ambient water temperature ≥ 8 milligrams per litre (5th percentile)
								Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Ideal category for full contact recreation.	≤ 130 counts/100ml (95th percentile)
							Habitat	Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	> 88% = A/B category
							Biota	Fish	FRAI score	Fish condition	> 88% = A/B category
								Invertebrates	MIRAI score	Macroinvertebrate condition	> 82% = B category

Table 5: Resource Quality Objectives for RIVERS in priority Resource Units in the Integrated Unit of Analysis B4 Lower Berg

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
B4 Lower Berg	III	G10J	B4-R09	Berg River	Bvi16	D	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a D category	Months	Oct	26.184	0.000	15.280	Nov	9.579	8.000	Jan	8.272	7.947	10.951	14.684	24.346	31.158	37.184	1.619																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
												Maintenance flows (million cubic metres)	Low	0.000	2.496	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Salts	Electrical conductivity (EC)	Salt concentrations need to be maintained at levels that do not adversely affect aquatic ecosystems	≤ 55 milliSiemens/metre (95th percentile)
								System variables	pH range Water temperature Dissolved oxygen	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	$6.5 \leq \text{pH} \leq 8.5$ (5th and 95th percentiles) $\pm 2^\circ\text{C}$ difference from ambient water temperature ≥ 6 milligrams litre (5th percentile)
								Toxins	Atrazine Endosulfan	Toxicity levels must not pose a threat to aquatic ecosystems.	≤ 0.079 milligrams per litre (95th percentile) ≤ 0.0013 milligrams per litre (95th percentile)
								Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for full contact recreation.	≤ 1000 counts/100ml (95th percentile)
								Geomorphology	GAI score - D50 VEGRAI level 3 score.	Geomorphological condition Sand particle size Vegetation condition	$> 68\% = \text{B/C category}$ $0.576 > \text{D50} > 0.349$ $> 42\% = \text{D category}$
									Exotic species Terrestrial woody species		No exotic plant species.
									Indigenous riparian woody species Non-woody indigenous species	Marginal zone cover abundance	No terrestrial woody species. Cover 30-50%.
									Reeds		Cover 30-50%.
									Exotic species		Cover 30-50%.
									Terrestrial woody species		Cover < 5%.
									Indigenous riparian woody species		Cover < 10%.
									Non-woody indigenous species	Lower zone cover abundance	Cover 50-75%.
									Reeds		Cover 5-10%.
									Exotic species		No reeds
									Terrestrial woody species		Cover < 10%.
									Indigenous riparian woody species	Upper zone cover abundance	Cover $\leq$ 15%.
									Non-woody indigenous species		Cover 30-50%.
									Non-woody indigenous species		Cover 30-50%.

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
B4 Lower Berg	III	G10K	B4-R10	Berg River	Bvii12	D	Quantity	Low flows High flows	FRAI score Exotic fish species MIRAI score SASS5 and ASPT score Number of families	Fish condition Indigenous species richness Macroinvertebrate condition SASS scores Diversity of invertebrate community	> 18% = F category No increase in the number of exotic fish present: <i>Cyprinus carpio</i> , <i>Oreochromis mossambicus</i> , <i>Tilapia sparrmanii</i> , <i>Micropterus punctulatus</i> , <i>Clarias gariepinus</i> and <i>Gambusia affinis</i> . > 42% = D category SASS5 score >80, ASPT ≥ 5.0 >/= 15 families, at an abundance of A to C.
B4 Lower Berg	III	G10K	B4-R10	Berg River	Bvii12	D	Quality	Nutrients Salts System variables Toxins Pathogens	Phosphate (PO ₄ -P) Total inorganic nitrogen (TIN) Electrical conductivity (EC) pH range Water temperature Dissolved oxygen Atrazine Endosulfan Escherichia coli	Nutrient levels must be maintained in the river at an mesotrophic condition. Salt concentrations need to be maintained at levels that do not adversely affect aquatic ecosystems pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health. Toxicity levels must not pose a threat to aquatic ecosystems. Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation.	≤ 0.075 milligrams/litre (50th percentile) ≤ 1.75 milligrams/litre (50th percentile) ≤ 55 milliSiemens/metre (95th percentile) 6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles) 2°C difference from ambient ≥ 6 milligrams litre (5th percentile) ≤ 0.079 milligrams per litre (95th percentile) ≤ 0.0013 milligrams per litre (95th percentile) ≤ 2500 counts/100ml (95th percentile)
B4 Lower Berg	III	G10K	B4-R10	Berg River	Bvii12	D	Habitat Habitat	Geomorphology Riparian vegetation	GAI score - D50 VEGRAI level 3 score. Exotic species	Geomorphological condition Sand particle size Vegetation condition Marginal zone cover	> 68% = B/C category 0.860 > D50 > 0.275 > 42% = D category No exotic plant species.

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Geomorphology Riparian vegetation	Terrestrial woody species Indigenous riparian woody species Non-woody indigenous species Reeds	abundance	No terrestrial woody species.
								Fish	FRAI score	Fish condition	Cover 30-50%
								Invertebrates Fish	Exotic fish species	Indigenous species richness	Cover 50-75%.
											Cover 15-25%.
											85% (B category)
											No increase in the number of exotic fish present: <i>Cyprinus carpio</i> , <i>Oreochromis mossambicus</i> , <i>Tilapia sparrmanii</i> , <i>Micropterus punctulatus</i> , <i>Clarias gariepinus</i> and <i>Gambusia affinis</i> .
									MIRAI score	Macroinvertebrate condition	81.4% (B/C category)
									SASS5 and ASPT score	SASS5 scores	SASS5 score >85, ASPT ≥ 4.2.
									Number of families	Diversity of invertebrate community	>/= 19 families, at an abundance of A to C.

Table 6: Resource Quality Objectives for RIVERS in priority Resource Units in the Integrated Unit of Analysis D10 Diep

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																								
D10 Diep	III	G21D	D10-R11	Diep River	Bv1	D	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a D category	Maintenance flows (million cubic metres)	Months	Low	High	0.026	0.003	0.000	0.000	0.015	0.021	0.043	0.090	0.130	0.157	0.106										
													0.079	0.053	0.029	0.020	0.017	0.000	0.000	0.116	0.294	0.120	0.473												
												Nutrient levels must be maintained in the river at a mesotrophic or better condition.												≤ 0.075 milligrams/litre (50th percentile)											
												Diep River is naturally saline and should be maintained in its current status.												≤ 1.75 milligrams/litre (50th percentile)											
												Quality	Salts	Electrical conductivity (EC)	pH range Water temperature Dissolved oxygen	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	6.5 ≥ pH ≤ 8.5 (5th and 95th percentiles)																		
																	2°C difference from ambient water temperature																		
							≥ 6 milligrams litre (5th percentile)																												

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
D10 Diep	III	G21D	D10-R12	Diep River	Biv6	D	Quantity	Toxins	Atrazine Endosulfan	Toxicity levels must not pose a threat to aquatic ecosystems.	≤ 0.079 milligrams per litre (95th percentile) ≤ 0.0013 milligrams per litre (95th percentile)
								Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation.	≤ 2500 counts/100ml (95th percentile)
								Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a D category	Months
											Low
											High
											Maintenance flows (million cubic metres)
							Quality	Nutrients	Phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)	River nutrient levels must be improved to eutrophic conditions.	≤ 0.125 milligrams/litre (50th percentile) ≤ 2.5 milligrams/litre (50th percentile)
									Electrical conductivity (EC)	Diep River is naturally saline and should be maintained in its current status.	≤ 350 millisiemens/metre (95th percentile)
								System variables	pH range	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles)
									Water temperature	2°C difference from ambient water temperature	
									Dissolved oxygen	≥ 6 milligrams litre (5th percentile)	
								Toxins	Atrazine Endosulfan	Toxicity levels must not pose a threat to aquatic ecosystems.	≤ 0.079 milligrams per litre (95th percentile) ≤ 0.0013 milligrams per litre (95th percentile)
							Habitat	Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation.	≤ 2500 counts/100ml (95th percentile)
								Geomorphology	GAI score	Geomorphological condition	> 22% = E category
								Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	> 18% = F category
							Biota	Fish	FRAI score	Fish condition	> 22% = E category
								Invertebrates	MIRAI score	Macroinvertebrate condition	> 22% = E category

Table 7: Resource Quality Objectives for RIVERS in priority Resource Units in the Integrated Unit of Analysis E11 Peninsula

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																								
E11 Peninsula	II	G22B	E11-R13	Hout Bay	Bviii6	D	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a D category	Months	Oct	0.037	0.132	0.071	0.038	0.000	0.029	0.000	0.026	0.000	0.025	0.037	0.121	0.070	0.142	0.221	0.252	0.204	0.188					
												Low	0.003	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
							Quality	Nutrients	Phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)	Nutrient levels must be maintained in the river in a eutrophic or better condition.	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	≤ 0.125 milligrams per litre (50th percentile)					≤ 2.50 milligrams per litre (50th percentile)									
												Low	0.037	0.132	0.071	0.038	0.000	0.029	0.000	0.026	0.000	0.025	0.037	0.121	0.070	0.142	0.221	0.252	0.204	0.188					
								Salts	Electrical conductivity (EC)	Salt concentrations need to be maintained at levels that do not adversely affect aquatic ecosystems	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	≤ 55 milliSiemens/metre (95th percentile)					6.5 ≥ pH ≤ 8.5 (5th and 95th percentiles)					2°C difference from ambient water temperature				
								System variables	pH range Water temperature Dissolved oxygen	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	≤ 4000 counts/100ml (95th percentile)					≥ 6 milligrams per litre (5th percentile)									
								Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for full contact recreation.	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	> 22% = E category					> 18% = E/F category									
Habitat	Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	> 42% = D category					> 42% = D category																
Biota	Fish	FRAI score	Fish condition	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204											> 42% = D category					> 42% = D category						
	Invertebrates	MIRAI score	Macroinvertebrate condition	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	> 42% = D category					> 42% = D category																
E11 Peninsula	II	G22A	E11-R14	Silvermine River	Bvii20	C	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a C category	Months	Oct	0.017											0.167	0.105	0.053	0.000	0.035	0.000	0.029	0.000	0.027	0.037	0.069	0.138
												Low	0.002	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
							Quality	Nutrients	Phosphate (PO ₄ -P)	Nutrient levels must be maintained in the river in a eutrophic or better condition.	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	≤ 0.075 milligrams/litre (50th percentile)					≤ 0.075 milligrams/litre (50th percentile)									
								Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	> 22% = E category					> 18% = E/F category									
							Biota	Fish	FRAI score	Fish condition	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	> 42% = D category					> 42% = D category									
							Invertebrates	Macroinvertebrate condition	Macroinvertebrate condition	Macroinvertebrate condition	Maintenance flows (million cubic metres)	High	0.188	0.094	0.543	0.302	0.142	0.221	0.252	0.204	> 42% = D category					> 42% = D category									

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Salts	Electrical conductivity (EC)	Salt concentrations need to be maintained at present day levels.	≤ 85 milliSiemens/metre (95th percentile)
								System variables	pH range Water temperature Dissolved oxygen	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles) 2°C difference from ambient water temperature ≥ 6 milligrams litre (5th percentile)
								Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in a Tolerable category for intermediate contact recreation. In the long term the aim should be to improve the river to an Acceptable, and then Ideal category for intermediate contact recreation.	≤ 4000 counts/100ml (95th percentile)
							Habitat	Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	> 38% = D/E category
							Biota	Fish	FRAI score	Fish condition	> 62% = C category

Table 9: Resource Quality Objectives for RIVERS in priority Resource Units in the Integrated Unit of Analysis D6 Eerste

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric													
D6 Eerste	III	G22F	D6-R16	Jonkershoek River	Biii6	C	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a C category	Months	Low	0.639	0.543	0.349	0.200	0.142	0.126	0.186	0.335	0.522	0.645	0.714	0.693
													High	0.245	0.067	0.000	0.000	0.000	0.000	0.454	0.747	1.052	0.206	0.412
													Maintenance flows (million cubic metres)											
													≤ 0.075 milligrams/litre (50th percentile)											
													≤ 1.75 milligrams/litre (50th percentile)											
													Nutrient levels must be maintained in the river at a mesotrophic or better condition.											
							Quality	Nutrients	Phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)	Electrical conductivity (EC)	Salt concentrations need to be maintained at present day levels.	≤ 55 milliSiemens/metre (95th percentile)												
												Salts												

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
D6 Eerste	III	G226	D6-R17	Klipies River	Biv8	D	Quantity	Low flows High flows	pH range Water temperature Dissolved oxygen	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles) ≥ 6 milligrams per litre (5th percentile)
							Quality	Nutrients	Ammonia Atrazine Endosulfan	Toxicity levels must not pose a threat to aquatic ecosystems.	≤ 0.073 milligrams per litre (95th percentile) ≤ 0.079 milligrams per litre (95th percentile) ≤ 0.0013 milligrams per litre (95th percentile)
							Habitat	Geomorphology Riparian vegetation	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation. In the long term the aim should be to improve the river to an Ideal category for intermediate contact recreation.	≤ 2500 counts/100ml (95th percentile)
							Biota	Fish	VEGRAI level 3 score.	Geomorphological condition Vegetation condition	> 62% = C category > 62% = C category
							Quantity	Maintenance low flows Maintenance high flows	Maintenance flows (million cubic metres)	Flows sufficient to maintain the river in a D category	≤ 0.125 milligrams/litre (50th percentile) ≤ 3.0 milligrams/litre (50th percentile)

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
									Endosulfan	Concentrations of waterborne pathogens should be maintained in a Tolerable category for intermediate contact recreation. In the long term the aim should be to improve the river to an Acceptable, and then Ideal category for intermediate contact recreation.	≤ 0.0013 milligrams per litre (95th percentile)
								Pathogens	Escherichia coli		≤ 4000 counts/100ml (95th percentile)
							Habitat	Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	> 22% = E category
							Biota	Fish	FRAI score	Fish condition	> 18% = D/E category
								Invertebrates	MIRAI score	Macroinvertebrate condition	> 62% = C category

Table 10: Resource Quality Objectives for RIVERS in priority Resource Units in the Integrated Unit of Analysis D7 Sir Lowrys

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																				
D7 Sir Lowry's	II	G22J	D7-R18	Lourens River	Bvii21	D	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a D category	Months	Oct	0.523	0.355	0.083	0.000	0.151	0.108	0.100	0.141	0.254	0.410	0.520	0.592	0.568						
											Maintenance flows (million cubic metres)	Low																			
								High	0.593			0.297	1.463	1.007	0.563	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
								≤ 0.075 milligrams/litre (50th percentile)																							
							≤ 1.75 milligrams/litre (50th percentile)																								
							Quality								Salts	Electrical conductivity (EC)	Salt concentrations need to be maintained at present day levels.	≤ 55 milliSiemens/metre (95th percentile)													
																pH range	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles)													
															System variables	Water temperature	Toxicity levels must not pose a	2°C difference from ambient water temperature													
																Dissolved oxygen		Toxicity levels must not pose a	≥ 6 milligrams litre (5th percentile)												
															Toxins	Ammonia	≤ 0.073 milligrams per litre (95th percentile)														

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric															
D7 Sir Lowry's	II	G22J	D7-R19	Sir Lowry's Pass River		C	Quantity	Low flows High flows	Maintenance low flows Maintenance high flows	Flows sufficient to maintain the river in a C category	Months	Low	High	0.380	0.959	0.000	0.000	0.000	0.204	0.186	0.257	0.459	0.755	0.984	1.141	1.145
											Maintenance flows (million cubic metres)															
											≤ 0.075 milligrams/litre (50th percentile)															
											≤ 1.75 milligrams/litre (50th percentile)															
											≤ 55 milliSiemens/metre (95th percentile)															
											6.5 ≤ pH ≤ 8.5 (5th and 95th percentiles)															
							Quality	Salts	Nutrients	Phosphate (PO ₄ P) Total inorganic nitrogen (TIN)	Electrical conductivity (EC)	Salt concentrations need to be maintained at present day levels.	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	Toxicity levels must not pose a threat to aquatic ecosystems.	2°C difference from ambient water temperature											
															≥ 6 milligrams litre (5th percentile)											
															≤ 0.073 milligrams per litre (95th percentile)											
															≤ 0.079 milligrams per litre (95th percentile)											
															≤ 0.0013 milligrams per litre (95th percentile)											

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																																								
D7 Sir Lowry's	II	G40A	D7-R20	Steenbras River	Bvii22	B/C	Quantity	Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation. In the long term the aim should be to improve the river to an Ideal category for intermediate contact recreation.	≤ 2500 counts/100ml (95th percentile)																																								
								Habitat	Riparian vegetation	VEGRAI level 3 score.	> 42% = D category																																								
								Biota	Fish	FRAI score	Fish condition	> 42% = D category																																							
									Invertebrates	MIRAI score	Macroinvertebrate condition	> 62% = C category																																							
								Quality	Low flows High flows	Maintenance low flows Maintenance high flows	Phosphate (PO ₄ P) Total inorganic nitrogen (TIN)	Nutrient levels must be maintained in the river at a oligotrophic condition.	Flows sufficient to maintain the river in a B/C category	Months	Low	Maintenance flows (million cubic metres)	0.000	0.427	Oct	0.000	0.323	Nov	0.000	0.235	Dec	0.000	0.180	Jan	0.000	0.149	Feb	0.000	0.144	Mar	0.000	0.173	Apr	0.077	0.247	May	0.384	Jun	0.307	0.506	Jul	0.307	0.582	Aug	0.077	0.502	Sep
														High	0.077	≤ 0.025 milligrams/litre (50th percentile)																																			
									Nutrients	Electrical conductivity (EC)				Salt concentrations need to be maintained at present day levels.	pH, temperature, and dissolved oxygen are important for the maintenance of ecosystem health.	5.0 ≤ pH ≤ 7.5 (5th and 95th percentiles) 2°C difference from ambient water temperature	≥ 6 milligrams litre (5th percentile)	Toxicity levels must not pose a threat to aquatic ecosystems.	Concentrations of waterborne pathogens should be maintained in an Acceptable category for full contact recreation.	≤ 165 counts/100ml (95th percentile)																															
																					Iron	≤ 0.1 milligrams per litre (95th percentile)																													
								Toxins	Manganese	Escherichia coli	Geomorphological condition	> 82% = B category																																							
													Pathogens	Geomorphology	GAI score	Geomorphological condition	> 82% = B category																																		

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Riparian vegetation	VEGRAI level 3 score.	Vegetation condition	> 78% = B/C category
							Biota	Fish	FRAI score	Fish condition	> 52% = D category
								Invertebrates	MIRAI score	Macroinvertebrate condition	> 92% = A category

Table 11: Resource Quality Objectives for ESTUARIES in priority Resource Units in the Integrated Unit of Analysis A1 Berg Estuary

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
A1 Berg Estuary	II	G10M	A1-E01	Berg (Groot) Estuary	Bxi1	C	Quantity	Surface flow	Flow	River inflow should never drop below $0.6 \text{ m}^3 \cdot \text{s}^{-1}$ and should not below $1 \text{ m}^3 \cdot \text{s}^{-1}$ for longer than 4 months; Flood frequency should not increase/decrease by more than 10% from 2004 baseline conditions	Oct 31.21 (46%) Nov 12.55 (36%) Dec 3.92 (25%) Jan 1.61 (19%) Feb 1.50 (23%) Mar 1.66 (20%) Apr 9.13 (36%) May 22.18 (26%) Jun 64.25 (42%) Jul 123.35 (61%) Aug 137.15 (68%) Sep 78.34 (63%) Annual 486.86 (52%)
								Nutrients	DIN	Inorganic nutrient concentrations not to exceed TPCs for macrophytes and microalgae	Estuary (low flows < $1 \text{ m}^3 \cdot \text{s}^{-1}$, summer): DIN <300 $\mu\text{g/l}$; DRP <100 $\mu\text{g/l}$ in Zones A and B, DIN <80 $\mu\text{g/l}$; DRP <30 $\mu\text{g/l}$ in Zones C and D Estuary (high flows > $5 \text{ m}^3 \cdot \text{s}^{-1}$, winter): DIN <800 $\mu\text{g/l}$; DRP <60 $\mu\text{g/l}$ in Zones A-D River inflow (< $1 \text{ m}^3 \cdot \text{s}^{-1}$, summer): DIN <80 $\mu\text{g/l}$; DRP <20 $\mu\text{g/l}$ River inflow (> $5 \text{ m}^3 \cdot \text{s}^{-1}$, winter): DIN <800 $\mu\text{g/l}$; DRP <60 $\mu\text{g/l}$
							Quality	Salinity	Salinity	Salinity distribution not to exceed TPCs for fish, invertebrates, macrophytes and microalgae	Salinity <20 for longer than 3 months at 20 km upstream from the mouth; Salinity <1 ppt above 40 km upstream of the mouth; Salinity of Salinity everywhere in estuary <35; Groundwater salinity on floodplain <45; TDS of river inflow <3500 mg/l "River inflow: 7 < pH < 8.5 Estuary: 7 < pH < 8.5" "River inflow: DO >4 mg/l Estuary DO >4 mg/l"
								System variables	Temperature	System variables not to exceed TPCs for biota	
									pH		
									Dissolved oxygen		
									Secchi depth		
								Pathogens	Enterococci	Concentrations of waterborne pathogens should be maintained in an Acceptable category for contact recreation	Zones A and B <1.0 m during low flow (< $1 \text{ m}^3 \cdot \text{s}^{-1}$) ≤185 Enterococci/100 ml) (90th percentile, hazen system)
							Habitat	Hydrodynamics	Mouth state	Habitat health adequate for	Permanently open

IUA Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Tidal variation	microalgae, macrophytes, invertebrates, fish, birds and recreational use	<10% change from present state
							Sediments	Sediment characteristics, Channel shape/size		Bathymetry and sediment MdØ change <10% from baseline
							Microalgae	Biomass and community composition of phytoplankton and benthic microalgae community	Phytoplankton biomass and composition suitable for invertebrates, fish, birds and recreational use	Blue-green algae <10% of phytoplankton cell counts, Benthic microphytobenthic < 40 mg/m ² chlorophyll a, The frequency of dinoflagellates < 5% of the total phytoplankton counts
									Macrophyte cover and composition suitable for invertebrates, fish, birds and recreational use	Maintain the present distribution (2003-2005) and abundance of the different plant community types and estuarine habitats (intertidal mudflats with <i>Zostera capensis</i> 206 ha, intertidal salt marsh 499 ha, open pan 1159 ha, halophytic floodplain 1521 ha, xeric floodplain 919.1 ha, reeds and sedges 586.6 ha and sedge pan 292.5 ha), Prevent an increase in mats of macroalgae in the lower intertidal reaches, Reduce the area covered by water hyacinth (<i>Eichornia crassipes</i>) in the upper reaches by 50% compared to the present state (2003-2005), Prevent an increase in size of the open pan dry areas (1159 ha in 2003-2005), Prevent a decrease in size of the sedge pan areas (293 ha in 2003-2005), <i>Juncus maritimus</i> , and waterblommetjies <i>Aponogeton distachyos</i> are present, Prevent the spread of invasive aliens in the riparian zone (e.g. <i>Acacia mearnsii</i> and <i>Eucalyptus camaldulensis</i>), Maintain intact reed and sedge stands along the banks of the estuary by ensuring that salinity is not greater than 20 ppt for 3 months at 20 km from the month during summer, Prevent an increase in bare ground in the halophytic and xeric floodplain habitats by maintaining the present-day flooding patterns
						Biota	Macrophytes	Extent, distribution and richness of macrophytes		
							Invertebrates	Macrofauna community composition, abundance and richness	Abundance and community composition of invertebrates suitable for fish, birds	Retain present species richness, distribution of species and mix (low species abundance, high dominance) in Zones A to the middle reaches of Zone C. One or two species will always be present at high densities compared to others (e.g. <i>Pseudodiaptomus hessei</i> , <i>Grandierella</i> sp.) in these Zones (A to C). Indicator species such as <i>Capitella capitata</i> , should not dominate benthic species at any site, <i>Callinassa kraussi</i> and <i>Upogebia africana</i> distribution patterns remain similar to present state.

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Fish	Fish community composition, abundance and richness	Abundance and community composition of fish community suitable for birds	Retain the full complement of estuarine resident (7 species) and estuary associated marine (5 species) present in the estuary with population sizes sufficient to ensure their persistence in perpetuity. Ensure that exotic freshwater species do not increase to levels where they can exclude any more indigenous species through predation or competitive interactions. Maintain recruitment of adult and juvenile fish at present levels. This requires maintaining sufficient flow for freshwater plume (temperature, salinity and olfactory gradient) entering the sea. This implies that there should be a significant number of 0-1-year-old fish and no missing year classes.
								Birds	Avifauna community composition, abundance and richness	Health avifauna community contributing to conservation of avifauna species in SA	Retain at least 90% of the baseline species richness, abundance and diversity of the bird community determined using regression slope based on a 3-year running average

Table 12: Resource Quality Objectives for ESTUARIES in priority Resource Units in the Integrated Unit of Analysis A2 Langebaan

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
A2 Langebaan	II	G10M	A2-E02	Langebaan	Bx13			Nutrients	NO ₃	Inorganic nutrient concentrations not to exceed TPCs for macrophytes and microalgae	NO ₃ < 1.3 mg.l ⁻¹
								Salinity	Salinity	Salinity distribution not to exceed TPCs for fish, invertebrates, macrophytes and microalgae	Salinity at the head of the lagoon < 40; Rest of the lagoon 34 < Salinity < 36
								System variables	Dissolved oxygen	System variables not to exceed TPCs for biota	> 4 mg.l ⁻¹
									Secchi depth		Secchi depth > 1 m
									Enterococci	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation	≤ 185 Enterococci/100 ml (90th percentile, hazen system)
								Pathogens	Escherichia coli		≤ 500 E. coli/100 ml (90th percentile, hazen system)

IUA Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
						Habitat	Hydrodynamics	Tidal amplitude	Habitat health adequate for microalgae, macrophytes, invertebrates, fish, birds and recreational use	Tidal amplitude should not change more than 10% from present state (2017)
							Sediments	Sediment characteristics, Channel shape/size	microalgae, macrophytes, invertebrates, fish, birds and recreational use	Bathymetry and sediment MdØ change <10% from baseline
						Biota	Microalgae	Biomass and community composition of phytoplankton and benthic microalgae community	Phytoplankton biomass and composition suitable for invertebrates, fish, birds and recreational use	Maintain low phytoplankton biomass (chlorophyll- a < 20 µg/l) and a diversity of phytoplankton groups.
							Macrophytes	Extent, distribution and richness of macrophytes	Macrophyte cover and composition suitable for invertebrates, fish, birds and recreational use	Maintain the distribution and area cover of macrophyte habitats particularly the salt marsh and seagrass. Maintain the large groundwater fed rush habitat.
							Invertebrates	Macrofauna community composition, abundance and richness	Abundance and community composition of Invertebrates suitable for fish, birds	In terms of Invertebrates Langebaan lagoon is currently in an A category. The invertebrate communities are in good health with species richness, abundances and composition scoring highly.
							Fish	Fish community composition, abundance and richness	Abundance and community composition of fish community suitable for birds	The fish community should include healthy populations of exploited fish species, specifically the harders, white stumpnose, blacktail, elf and smooth hound shark juveniles should all be present in beach seine net sampling surveys (at least 10 hauls in 3 different sites) of the near-shore areas. Adults of these species should remain the main components in the catches of line and net fisheries in the lagoon, and catch rates should remain stable or increase.
									Health avifauna community contributing to conservation of avifauna species in SA	Retain at least 90% of the baseline species richness, abundance and diversity of the bird community determined using regression slope based on a 3-year running average.
							Birds	Avifauna community composition, abundance and richness		

Table 13: Resource Quality Objectives for ESTUARIES in priority Resource Units in the Integrated Unit of Analysis D10 Diep

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																															
D10 Diep	III	G21F	D10-E03	Rietvlei/Diep	Bviii5	D	Quantity	Surface flow	Flow	Freshwater inflow adequate to maintain water quality and habitat suitable for flora and fauna	Months	Oct	80 %	Nov	80 %	Dec	80 %	Jan	93 %	Feb	100 %	Mar	100 %	Apr	80 %	May	80 %	Jun	80 %	Jul	80 %	Aug	80 %	Sep	80 %	Annual						
											MMR/MAR (% Nat)																															
											River inflow: <800 µg.l ⁻¹																															
							Quality	Nutrients	DIN	Inorganic nutrient concentrations not to exceed TPCs for macrophytes and microalgae		Lower estuary (Milnerton lagoon): <1000 µg.l ⁻¹																														
											DIP	River inflow: <60 µg.l ⁻¹																														
									Salinity	Salinity distribution not to exceed TPCs for fish, invertebrates, macrophytes and microalgae		Average salinity in lower estuary (Milnerton lagoon) = 20, maximum = 35																														
							System variables	System variables	Dissolved oxygen	System variables (temperature, pH, dissolved oxygen, suspended solids and turbidity) not to exceed TPCs for biota		>4 mg.l ⁻¹																														
											Enterococci	≤185 Enterococci/100 ml (90th percentile, hazen system)																														
								Pathogens	Escherichia coli	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation		≤500 E. coli/100 ml (90th percentile, hazen system)																														
							Habitat	Hydrodynamics	Mouth state	Habitat health adequate for microalgae, macrophytes, invertebrates, fish, birds and recreational use		Permanently open																														
											Tidal variation	<10% change from present state																														
								Sediments	Sediment characteristics, Channel shape/size	Bathymetry and sediment MdØ change <10% from baseline																																
							Biota	Microalgae	Biomass and community composition of phytoplankton and benthic microalgae community	Phytoplankton biomass and composition suitable for invertebrates, fish, birds and recreational use		Maintain low phytoplankton biomass (chlorophyll- a < 50 µg/l) and a diversity of phytoplankton groups.																														

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Macrophytes	Extent, distribution and richness of macrophytes	Macrophyte cover and composition suitable for invertebrates, fish, birds and recreational use	Maintain the distribution and area cover of macrophyte habitats particularly the salt marsh
								Invertebrates	Macrofauna community composition, abundance and richness	Abundance and community composition of invertebrates suitable for fish, birds	Restore and maintain species richness, distribution of species and mix (low species abundance, high dominance); Indicator species such as <i>Capitella capitata</i> , should not dominate benthic species at any site; <i>Callinassa kraussi</i> and <i>Upogebia africana</i> distribution patterns similar to reference state.
								Fish	Fish community composition, abundance and richness	Abundance and community composition of fish community suitable for birds	Restore and maintain the full complement of estuarine resident and estuary associated marine present in the estuary with population sizes sufficient to ensure their persistence in perpetuity; Ensure that exotic freshwater species do not increase to levels where they can exclude any more indigenous species through predation or competitive interactions; Maintain recruitment of adult and juvenile fish at present levels.
								Birds	Avifauna community composition, abundance and richness	Health avifauna community contributing to conservation of avifauna species in SA	Retain at least 90% of the baseline species richness, abundance and diversity of the bird community determined using regression slope based on a 3-year running average.

Table 14: Resource Quality Objectives for ESTUARIES in priority Resource Units in the Integrated Unit of Analysis E11 Peninsula

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
E11 Peninsula	II	G22A	E11-E04	Wildevoelvi	Bxi14	D	Quantity	Surface flow	Flow	Freshwater inflow does not exceed requirements for maintaining water quality and habitat suitable for flora and fauna	Months	120 %	Oct	120 %	Nov	120 %	Dec	120 %	Jan	120 %	Feb	120 %	Mar	120 %	Apr	120 %	May	120 %	Jun	120 %	Jul	120 %	Aug	120 %	Sep	120 %	Annual	120 %																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																														
											MMR/MAR (% Nat)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									</

IUA Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
							Salinity	Salinity	Salinity distribution not to exceed TPCs for fish, invertebrates, macrophytes and microalgae	Average salinity in lower estuary (backshore lagoon) >10, maximum = 35, average salinity in Wildevoei vlei > 2
							System variables	Dissolved oxygen	System variables not to exceed TPCs for biota	>4 mg.l ⁻¹
								Enterococci	Concentrations of waterborne pathogens should be maintained in an Acceptable category for full contact recreation	≤185 Enterococci/100 ml (90th percentile, hazen system)
							Pathogens	Escherichia coli		≤500 E. coli/100 ml (90th percentile, hazen system)
							Hydrodynamics	Mouth state	Habitat health adequate for microalgae, macrophytes, invertebrates, fish, birds and recreational use	Mouth should remain open >70% of the time
								Tidal variation		<10% change from present state
							Sediments	Sediment characteristics, Channel shape/size		Bathymetry and sediment MqØ change <10% from baseline
								Biomass and community composition of phytoplankton and benthic microalgae	Phytoplankton biomass and composition suitable for invertebrates, fish, birds and recreational use	Improvement from current hypereutrophic state where toxic cyanobacteria are common and flow to the sea
								Extent, distribution and richness of macrophytes	Macrophyte cover and composition suitable for invertebrates, fish, birds and recreational use	Retain present species richness, distribution of species and mix (low species abundance, high dominance); Maintain the fringing vegetation around the vleis as this is important for bank stabilisation and nutrient uptake; Improve connectivity between the sea, channel and lower vlei; Control the spread of invasive floating aquatic macrophyte species present in the vleis e.g. water fern.
							Macrophytes			
							Invertebrates	Macrofauna community composition, abundance and richness	Abundance and community composition of Invertebrates suitable for fish, birds	Move from a D category to a C category. The estuary should have a viable population of <i>Callichirus kraussi</i> in the backwater lagoon (10/m2). In addition, the invertebrate community should include 2 other estuarine species in the canal. At least three marine invertebrate species present near the mouth.

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Fish	Fish community composition, abundance and richness	Abundance and community composition of fish community suitable for birds	Maintain fish assemblage that includes at least two species of mullet, <i>Liza richardsonii</i> and either/both <i>Mugil cephalus</i> and <i>Pseudomyxus capensis</i> . Substantial seasonal fluctuations in abundance of these mullet species are expected to occur, but mullet should remain more abundant than the alien freshwater species currently inhabiting the vleis.
								Birds	Avifauna community composition, abundance and richness	Health avifauna community contributing to conservation of avifauna species in SA	Retain at least 90% of the baseline species richness, abundance and diversity of the bird community determined using regression slope based on a 3-year running average.

Table 15: Resource Quality Objectives for ESTUARIES in priority Resource Units in the Integrated Unit of Analysis E12 Cape Flats

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
E12 Cape Flats	III	G22K	E12-E05	Zandvlei	Bx19	D	Quantity	Surface flow	Flow	Freshwater inflow adequate to maintain water quality and habitat suitable for flora and fauna.	Months
											Oct
											Nov
											Dec
											Jan
											Feb
											Mar
											Apr
											May
											Jun
											Jul
											Aug
											Sep
											Annual
							Quality	Nutrients	DIN	Inorganic nutrient concentrations not to exceed TPCs for macrophytes and microalgae	River inflow: <1000 µg.l-1
											Estuary: <150 µg.l-1
									DIP	TPCs for macrophytes and microalgae	River inflow: <300 µg.l-1
											Estuary: <100 µg.l-1
							Quality	Salinity	Salinity	Salinity distribution not to exceed TPCs for fish, invertebrates, macrophytes and microalgae	15 < Average salinity <35
							Quality	System variables	Dissolved oxygen	System variables not to exceed TPCs for biota	>4 mg.l-1
							Quality	Pathogens	Enterococci	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation.	≤185 Enterococci/100 ml (90th percentile, hazen system)
											≤500 E. coli/100 ml (90th percentile, hazen system)
							Habitat	Hydrodynamics	Mouth state	Habitat health adequate for contact recreation.	Mouth should remain open >20% of the time

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric	
E12 Cape Flats	III	G22K	E12-E05	Zeekoevlei	Bxi20	D	Quantity	Surface flow	Flow	Freshwater inflow adequate to maintain water quality and habitat suitable for flora and fauna	Months	120 %
											Qct	120 %
											Nov	120 %
											Dec	120 %
											Jan	120 %
											Feb	120 %
											Mar	120 %
											Apr	120 %
											May	120 %
											Jun	120 %
											Jul	120 %
											Aug	120 %
											Sep	120 %
											Annual	
Retain at least 90% of the baseline species richness, abundance and diversity of the bird community determined using regression slope based on a 3-year running average.												
Health avifauna community contributing to conservation of avifauna species in SA												
Birds												
Avifauna community composition, abundance and richness												
Fish												
Fish community composition, abundance and richness												
Invertebrates												
Macrofauna community composition, abundance and richness												
Abundance and community composition of Invertebrates suitable for fish, birds												
Biota												
Macrophytes												
Extent, distribution and richness of macrophytes												
Macrophyte cover and composition suitable for invertebrates, fish, birds and recreational use												
Microalgae												
Biomass and community composition of phytoplankton and invertebrates, fish, birds and benthic microalgae recreational use												
Phytoplankton biomass and composition suitable for invertebrates, fish, birds and recreational use												
Maintain low phytoplankton biomass (chlorophyll- a < 20 µg/l) and a diversity of phytoplankton groups.												
Maintain and/or restore distribution and area cover of macrophyte habitats particularly salt marsh												
Restore and maintain species richness, distribution of species and mix (low species abundance, high dominance); Indicator species such as <i>Capitella capitata</i> , should not dominate benthic species at any site; <i>Callianassa kraussi</i> and <i>Upogebia africana</i> distribution patterns similar to reference state.												
Restore and maintain the full complement of estuarine resident and estuary associated marine present in the estuary with population sizes sufficient to ensure their persistence in perpetuity; Ensure that exotic freshwater species do not increase to levels where they can exclude any more indigenous species through predation or competitive interactions; Maintain recruitment of adult and juvenile fish at present levels.												
Retain at least 90% of the baseline species richness, abundance and diversity of the bird community determined using regression slope based on a 3-year running average.												

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
										concentrations not to exceed TPCs for macrophytes and microalgae	Lower estuary: <1000 µg.l ⁻¹ River inflow: <500 µg.l ⁻¹ Lower estuary: <500 µg.l ⁻¹
								Salinity	Salinity	Salinity distribution not to exceed TPCs for fish, invertebrates, macrophytes and microalgae	Average salinity in lower >10, maximum = 35
								System variables	Dissolved oxygen	System variables (temperature, pH, turbidity, dissolved oxygen, suspended solids and turbidity) not to exceed TPCs for biota	>4 mg.l ⁻¹
								Pathogens	Enterococci	Concentrations of waterborne pathogens should be maintained in an Acceptable category for intermediate contact recreation	≤185 Enterococci/100 ml (90th percentile, hazen system)
									Escherichia coli		≤500 E. coli/100 ml (90th percentile, hazen system)
							Habitat	Hydrodynamics	Mouth state	Habitat health adequate for microalgae, macrophytes, invertebrates, fish, birds and recreational use	Mouth should remain open >30% of the time
								Microalgae	Biomass and community composition of phytoplankton and benthic microalgae community	Phytoplankton biomass and composition suitable for invertebrates, fish, birds and recreational use	Phytoplankton biomass (measured as chlorophyll-a) <100 µg/l and a diversity of phytoplankton groups.
							Biota	Macrophytes	Extent, distribution and richness of macrophytes	Macrophyte cover and composition suitable for invertebrates, fish, birds and recreational use	Maintain and/or restore distribution and area cover of macrophyte habitats particularly salt marsh
								Invertebrates	Macrofauna community composition, abundance and richness	Restore and maintain species richness, distribution of species and mix (low species abundance, high dominance); Indicator species such as <i>Capitella capitata</i> , should not dominate benthic species at any site; <i>Callinassa kraussi</i> and <i>Upogebia africana</i> distribution patterns similar to reference state.	

IUA Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
							Fish	Fish community composition, abundance and richness	Abundance and community composition of fish community suitable for birds	Restore and maintain the full complement of estuarine resident and estuary associated marine present in the estuary with population sizes sufficient to ensure their persistence in perpetuity; Ensure that exotic freshwater species do not increase to levels where they can exclude any more indigenous species through predation or competitive interactions; Maintain recruitment of adult and juvenile fish at present levels.
							Birds	Avifauna community composition, abundance and richness	Health avifauna community contributing to conservation of avifauna species in SA	Retain at least 90% of the baseline species richness, abundance and diversity of the bird community determined using regression slope based on a 3-year running average.

Table 16: Resource Quality Objectives for ESTUARIES in priority Resource Units in the Integrated Unit of Analysis D6 Eerste

IUA Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																					
D6 Eerste	G22H	D6-E06	Eerste Estuary	Bxi3	D	Quantity	Surface flow	Flow	Freshwater inflow adequate to maintain water quality and habitat suitable for flora and fauna	Months	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Annual								
											120 %	120 %	120 %	120 %	120 %	120 %	120 %	120 %	120 %	120 %	120 %	120 %	120 %	120 %	120 %						
										MMR/MAR (% Nat)																					

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
									Tidal variation	microalgae, macrophytes, invertebrates, fish, birds and recreational use	<10% change from present state
								Microalgae	Biomass and community composition of phytoplankton and benthic microalgae community	Phytoplankton biomass and composition suitable for invertebrates, fish, birds and recreational use	Maintain low phytoplankton biomass (chlorophyll- <i>a</i> < 20 µg/l) and a diversity of phytoplankton groups.
								Macrophytes	Extent, distribution and richness of macrophytes	Macrophyte cover and composition suitable for invertebrates, fish, birds and recreational use	Restore and maintain the distribution and area cover of macrophyte habitats particularly salt marsh
								Invertebrates	Macrofauna community composition, abundance and richness	Abundance and community composition of Invertebrates suitable for fish, birds	Restore and maintain species richness, distribution of species and mix (low species abundance, high dominance); Indicator species such as <i>Capitella capitata</i> , should not dominate benthic species at any site; <i>Callinassa kraussi</i> and <i>Upogebia africana</i> distribution patterns similar to reference state.
								Fish	Fish community composition, abundance and richness	Abundance and community composition of fish community suitable for birds	Restore and maintain the full complement of estuarine resident and estuary associated marine present in the estuary with population sizes sufficient to ensure their persistence in perpetuity; Ensure that exotic freshwater species do not increase to levels where they can exclude any more indigenous species through predation or competitive interactions; Maintain recruitment of adult and juvenile fish at present levels.
								Birds	Avifauna community composition, abundance and richness	Health avifauna community contributing to conservation of avifauna species in SA	Retain at least 90% of the baseline species richness, abundance and diversity of the bird community determined using regression slope based on a 3-year running average.

Table 17: Resource Quality Objectives for ESTUARIES in priority Resource Units in the Integrated Unit of Analysis D7 Sir Lowry's

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric																																
D7 Sir Lowry's	II	G221	D7-E07	Lourens Estuary	Bxi4	D	Quantity	Surface flow	Flow	Freshwater inflow adequate to maintain water quality and habitat suitable for flora and fauna	Months	Oct	83 %	Nov	56 %	Dec	27 %	Jan	16 %	Feb	10 %	Mar	18 %	Apr	35 %	May	49 %	Jun	78 %	Jul	89 %	Aug	90 %	Sep	88 %	Annual	76 %						
											MMR/MAR (% Nat)																																
												River inflow: <350 µg.l ⁻¹																															
												Lower estuary: <300 µg.l ⁻¹																															
							Quality	Nutrients	DIN	Inorganic nutrient concentrations not to exceed TPCs for macrophytes and microalgae	River inflow: <80 µg.l ⁻¹																																
									DIP	TPCs for macrophytes and microalgae	Lower estuary: <80 µg.l ⁻¹																																
								Salinity	Salinity	Salinity distribution not to exceed TPCs for fish, invertebrates, macrophytes and microalgae	Average salinity in lower estuary >15, maximum = 35																																
							Habitat	Hydrodynamics	Mouth state	Habitat health adequate for microalgae, macrophytes, invertebrates, fish, birds and recreational use	Permanently open																																
												Tidal variation	<10% change from present state																														
								Sediments	Sediment characteristics, Channel shape/size	Bathymetry and sediment MddØ change <10% from baseline																																	
							Biota	Microalgae	Biomass and community composition of phytoplankton and benthic microalgae recreational use	Phytoplankton biomass and composition suitable for invertebrates, fish, birds and recreational use	Maintain low phytoplankton biomass (chlorophyll- a < 20 µg/l) and a diversity of phytoplankton groups.																																
								Macrophytes	Macrophyte cover and composition suitable for invertebrates, fish, birds and recreational use	Restore and maintain the distribution and area cover of macrophyte habitats particularly salt marsh																																	

IUA	Class	Quaternary Catchment	RU	Resource Name	Biophysical Node Name	TEC	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
								Invertebrates	Macrofauna community composition, abundance and richness	Abundance and community composition of Invertebrates suitable for fish, birds	Restore and maintain species richness, distribution of species and mix (low species abundance, high dominance); Indicator species such as <i>Capitella capitata</i> , should not dominate benthic species at any site; Callianassa kraussi and Upogebia africana distribution patterns similar to reference state.
								Fish	Fish community composition, abundance and richness	Abundance and community composition of fish community suitable for birds	Restore and maintain the full complement of estuarine resident and estuary associated marine present in the estuary with population sizes sufficient to ensure their persistence in perpetuity; Ensure that exotic freshwater species do not increase to levels where they can exclude any more indigenous species through predation or competitive interactions; Maintain recruitment of adult and juvenile fish at present levels.
								Birds	Avifauna community composition, abundance and richness	Health avifauna community contributing to conservation of avifauna species in SA	Retain at least 90% of the baseline species richness, abundance and diversity of the bird community determined using regression slope based on a 3-year running average.

Table 18: Resource Quality Objectives for DAMS in priority Resource Units in the Berg Catchment

UA Class	Quaternary Catchment	RU	Resource Name	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric														
D8 Upper Berg	G10A	D8-D01	Berg Dam	Quantity	Low flows	Dam level Flow releases: Berg EWR1 in G10A nMAR = 141.68 million m3/a pMAR: 126.00 million m3/a REC = C category	During the dry season dam levels must be sufficient for releases for irrigation and human use and protection of ecosystem function downstream. Water intake temperature to be managed.	Months	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Annual	
						Maintenance low flows (million cubic metres)	2.143	1.293	1.071	0.803	0.726	0.000	0.000	0.000	4.285	4.285	4.285	3.888	29.177			
				Quality	High flows		During the wet season high flow ecological releases are made according to the decision-support system.	Maintenance high flows (million cubic metres)	0.000	0.544	0.544	0.000	0.000	0.000	0.000	0.778	0.000	4.666	10.109	0.000	0.000	11.839
					Nutrients	Ortho-phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)1	The system must be maintained in a mesotrophic (moderately enriched) state or better to protect against nuisance algal blooms and excessive water treatment costs.	≤ 0.015 milligrams/litre (50 th percentile)														
				Quality			≤ 0.07 milligrams/litre (50 th percentile)															

IUA Class	Quaternary Catchment	RU	Resource Name	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
D8 Upper Berg	G10B	D8-D02	Wemmershoek Dam	Quantity	Salts	Electrical conductivity	Salt levels must be maintained at concentrations where they do not impact negatively on the ecosystem, are maintained in an Ideal category for domestic and irrigation water supply.	≤ 30 millisiemens/metre (95 th percentile)
					System variables	pH	The water in the dam is naturally acidic and it should be maintained within the historical range.	$5.5 \geq \text{pH} \leq 7.5$ (5 th and 95 th percentiles)
					Pathogens	E coli	The dam must be maintained in a state that is in an Ideal category for full contact recreation to protect its domestic water supply purpose.	≤ 130 counts/100ml (95 th percentile)
				Quality	Low flows	Dam levels	Dam levels must be sufficient for urban and industrial use water supply, and to supply some irrigators.	% of dam volume. No EWR site
					Nutrients	Ortho-phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)	The reservoir is currently in a Natural state and should be kept in an oligotrophic state. for supply to the City of Cape Town and Paarl. As a key domestic water supply reservoir this status should be maintained and protected.	≤ 0.005 milligrams/litre (50 th percentile)
						Ortho-phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)	The reservoir is currently in a Natural state and should be kept in an oligotrophic state. for supply to the City of Cape Town and Paarl. As a key domestic water supply reservoir this status should be maintained and protected.	≤ 0.50 milligrams/litre (50 th percentile)
B4 Lower Berg	G10F	B4-D03	Voelvlei Dam	Quantity	Low flows	Dam levels	Dam levels must be sufficient for urban and industrial use water supply via the two WTWs, and releases to Berg River for human and irrigation use.	% of dam volume. No EWR site
				Quality	Nutrients	Ortho-phosphate (PO ₄ -P)	The reservoir is currently in an Eutrophic state and should be	≤ 0.025 milligrams/litre (50 th percentile)

IUA Class	Quaternary Catchment	RU	Resource Name	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric
B4 Lower Berg	G10K	B4-D04	Misverstand Weir	Quantity		Total inorganic nitrogen (TIN)	Improved to a mesotrophic state or better to protect the water supply to the City of Cape Town and Swartland towns against harmful algal blooms and taste & odour problems in treated domestic water.	≤ 0.70 milligrams/litre (50 th percentile)
						Electrical conductivity	Salt levels must be maintained at concentrations where they do not impact negatively on the ecosystem, and are in an Ideal category for domestic water use and for irrigation water use.	≤ 30 milliSiemens/metre (95 th percentile)
					Pathogens	E coli, Faecal coliforms	The system must be maintained in a state that is in an Acceptable category for intermediate contact recreation	≤ 2000 counts/100ml (95 th percentile)
					Low flows	Dam levels	Water levels in the weir must be sufficient for supply for human use via the Withoogte WTW.	% of dam volume
					Nutrients	Ortho-phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)	The reservoir is currently in a Eutrophic state and should be in the short term be maintained in its current state or better. The long-term objective should be to improve the nutrient status to a mesotrophic state or better to protect the water supply to the West Coast towns.	≤ 0.025 milligrams/litre (50 th percentile)
						Ortho-phosphate (PO ₄ -P) Total inorganic nitrogen (TIN)		≤ 2.5 milligrams/litre (50 th percentile)
				Quality	Salts	Electrical conductivity	Salt levels must be maintained at concentrations where they do not impact negatively on the ecosystem, and are in an Ideal category for domestic and industrial water use, and for irrigation water use.	≤ 70 milliSiemens/metre (95 th percentile)
					Pathogens	E. coli	The reservoir must be maintained in a state that is safe for domestic water use (with treatment) and for intermediate contact recreation as the dam is a popular recreation venue.	≤ 1000 counts/100 ml (95 th percentile)
						Faecal coliforms		≤ 1000 counts/100 ml (95 th percentile)

UA Class	Quaternary Catchment	RU	Resource Name	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric					
D7 Sir Lowry's	G40A	D7-D05	Upper Steenbras Dam	Quantity	Low flows	Dam levels	Dam levels must be sufficient for releases to the Lower Steenbras Dam for urban and industrial use and protection of ecosystem functioning downstream of the Lower Steenbras Dam, hydropower energy generation via the Steenbras Pumped Storage Scheme as well as for water supply to the Western Cape Water Supply System (City of Cape Town) via the Faure WTW.	% of dam volume					
				Quality	Nutrients	Ortho-phosphate (PO ₄ -P)	The system must be maintained in a mesotrophic state or better.	≤ 0.015 milligrams/litre (50 th percentile)					
						Total inorganic nitrogen (TIN)							
						Ortho-phosphate (PO ₄ -P)		≤ 0.07 milligrams/litre (50 th percentile)					
						Total inorganic nitrogen (TIN)							
		Salts	Electrical conductivity	Salt levels must be maintained at concentrations where they do not impact negatively on the ecosystem, and are in an Ideal category for domestic and industrial water use, and for hydropower generation.	≤ 30 milliSiemens/metre (95 th percentile)								
		Pathogens	E. coli	The system must be maintained in a state that is safe for municipal use (with treatment).	≤ 130 counts/100 ml (95 th percentile)								
					Faecal coliforms	≤ 130 counts/100 ml (95 th percentile)							
D7 Sir Lowry's	G40A	D7-D06	Lower Steenbras Dam	Quantity	Low flows	Dam level Spills from dam. Flow releases: Berg EWR8 in G40A below Lower Steenbras Dam nMAR = 54.88 million m3/a	Dam levels must remain sufficient to provide for supply to the Western Cape Water Supply System (City of Cape Town) via the Steenbras WTW, and low flows to the lower Steenbras River and estuary for the protection of ecosystem functioning downstream.	Months					
								Oct	0.427				
								Nov	0.323				
								Dec	0.235				
								Jan	0.180				
												Feb	0.149
												Mar	0.144
												Apr	0.173
												May	0.247
												Jun	0.384
								Jul	0.506				
								Aug	0.582				
								Sep	0.502				
								Annual	3.852				

Quaternary Catchment	RU	Resource Name	Component	Sub-component	Indicator	RQO Narrative	RQO Numeric										
			High flows			High flow ecological releases should be made during the wet season to meet flood requirements, but within the constraints of the existing outlet structure, and utilising spills where possible.	Maintenance high flows (million cubic metres)	0.000	0.000	0.000	0.000	0.077	0.077	0.307	0.307	0.077	0.845
			Nutrients			Ortho-phosphate (PO ₄ -P)	The reservoir must be maintained in a mesotrophic state or better.	≤ 0.015 milligrams/litre (50 th percentile)									
						Total inorganic nitrogen (TIN)											
						Electrical conductivity											
Quality				Salt levels must be maintained at concentrations where they do not impact negatively on the ecosystem, and are in an ideal category for domestic and industrial water use.	≤ 30 milliSiemens/metre (95 th percentile)												
			E. coli	≤ 130 counts/100 ml (95 th percentile)													
			Faecal coliforms	≤ 130 counts/100 ml (95 th percentile)													

Table 19: Resource Quality Objectives for GROUNDWATER in priority Resource Units in the Berg Catchment

IUA Class	Quaternary Catchment	RU	Resource Name	Component	Sub Component	Indicator/ Measure	RQO Narrative	RQO Numeric
D8 Upper Berg	G10A	4-Paarl-Upper Berg	Groundwater (all)	Quantity	Abstraction	Seasonal abstraction: water level recovers from abstraction impact during wet season, under consideration of climate change and drought cycles. Permanent abstraction: water level decline stabilises under consideration of aquifer response time.	Groundwater use should be sustainable for all users and the environment	n/a
						Compliance with the low flow requirements in the river (as per riverine RQO)	Maintain (groundwater component of) the low flow requirements in the river	Maintenance low flow requirements: 29.177 Mm3/a (34.39 %MAR) at G1H076 (Bvii13); 27.421 Mm3/a (19.35 %MAR) at G1H077 (Bviii1)
					Nutrients	NO ₃ (as N)	Groundwater should be fit for domestic use after treatment;	< 3.3 mg/l
					Salts	EC	and groundwater quality shall not show a deteriorating trend from natural background	< 70 mS/m
					System variable	pH		5.2 – 8.4
					Pathogens	E-coli		0 counts / 100 ml
					Pathogens	Total Coliform		<10 counts / 100ml
				Quantity	Discharge	Relative water levels between groundwater and surface water (in mamsl)	The natural gradient between groundwater and surface water should be maintained	n/a
					Discharge	Buffer zones	No groundwater abstraction around wetland and river FEPAs in accordance with the implementation manual for FEPAs.	250m
				Quantity	Nutrients	NO ₃ (as N)	Groundwater should be fit for domestic use after treatment;	< 3.3 mg/l
C5 Berg Tributaries	G10B	4-Paarl-Upper Berg	Groundwater (all)	Quantity	Salts	EC	and groundwater quality shall not show a deteriorating trend from natural background	< 70 mS/m
					System variable	pH		5.2 – 8.4
					Pathogens	E-coli		0 counts / 100 ml
					Pathogens	Total Coliform		<10 counts / 100ml
				Quantity	Abstraction	Seasonal abstraction: water level recovers from abstraction impact during wet season, under consideration of climate change and drought cycles. Permanent abstraction: water level decline stabilises under consideration of aquifer response time.	Groundwater use should be sustainable for all users and the environment	n/a
					Abstraction			
				Quantity	Discharge	Relative water levels between groundwater and surface water (in mamsl)	The natural gradient between groundwater and surface water should be maintained	n/a
					Discharge	Buffer zones	No groundwater abstraction around wetland and river FEPAs in accordance with the implementation manual for FEPAs.	250m
				Quantity	Nutrients	NO ₃ (as N)	Groundwater should be fit for domestic use after treatment;	< 3.3 mg/l
					Salts	EC	and groundwater quality shall not show a deteriorating trend from natural background	< 70 mS/m

IUA	Class	Quaternary Catchment	RU	Resource Name	Component	Sub Component	Indicator/ Measure	RQO Narrative	RQO Numeric
B4 Lower Berg	III	G10J	6-24 Rivers		Quantity	Discharge	Buffer zones	No groundwater abstraction around wetland and river FEPAs in accordance with the implementation manual for FEPAs.	250m
						Pathogens	E-coli	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	0 counts / 100 ml
						Pathogens	Total Coliform		<10 counts / 100ml
					Quality	Nutrients	NO3 (as N)	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	n/a
						System variable	pH		n/a
						Salts	EC		n/a
				Groundwater (all)	Quantity	Discharge	Relative water levels between groundwater and surface water (in mamsl)	The natural gradient between groundwater and surface water should be maintained	n/a
							Buffer zones	No groundwater abstraction around wetland and river FEPAs in accordance with the implementation manual for FEPAs.	250m
							Compliance with the low flow requirements in the river (as per riverine RQO)	Maintain (groundwater component of) the low flow requirements in the river	Maintenance low flow requirements: 114.338 Mm ³ /a (13.28 %MAR) at G1H013 (Bvili6)
					Quality	System variable	pH		5.2 – 8.1
						Pathogens	E-coli	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	0 counts / 100 ml
						Pathogens	Total Coliform		<10 counts / 100ml
				Groundwater (Cenozoic coastal sand)	Quality	Nutrients	NO3 (as N)		< 6.9 mg/l
						Salts	EC		< 942 mS/m
				Groundwater (Basement)	Quality	Nutrients	NO3 (as N)		<11.0 mg/l
						Salts	EC		< 875 mS/m

IUA	Class	Quaternary Catchment	RU	Resource Name	Component	Sub Component	Indicator/ Measure	RQO Narrative	RQO Numeric
A1 Berg Estuary and A2 Langebaan	II	G10M	8-West Coast		Quantity	Abstraction	Seasonal abstraction: water level recovers from abstraction impact during wet season, under consideration of climate change and drought cycles. Permanent abstraction: water level decline stabilises under consideration of aquifer response time.	Groundwater use should be sustainable for all users and the environment	n/a
						Groundwater level	Water level	Minimum water level in abstraction boreholes within 2.5km from the ocean to avoid saline intrusion	>1 mamsl
						Discharge	Relative water levels between groundwater and surface water (in mamsl)	The natural gradient between groundwater and surface water should be maintained	n/a
							Buffer zones	No groundwater abstraction around wetland and river FEPAs in accordance with the implementation manual for FEPAs.	250m
							Compliance with the groundwater flow requirements to the Langebaan Lagoon	Compliance to the groundwater flow requirements to the Langebaan Lagoon, as per estuary RQO requirement	Groundwater inflow not <10% of present day (2017) rate
							Compliance with the groundwater flow requirements to the Langebaan Lagoon	Compliance to the groundwater flow requirements to the Langebaan Lagoon, as per estuary RQO requirement	Ground water level not <10% below present day (2017) level
				Groundwater (Cenozoic coastal sand)	Quality	Nutrients	NO3 (as N)	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 11.0 mg/l
						System variable	pH		7.1 - 8.4
						Salts	EC		< 520 mS/m
				Groundwater	Quality	Nutrients	NO3 (as N)	Groundwater should be fit for	< 11.0 mg/l

IUA	Class	Quaternary Catchment	RU	Resource Name	Component	Sub Component	Indicator/ Measure	RQO Narrative	RQO Numeric
				(Basement)		Salts	EC	domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 1571 mS/m
				Groundwater (all)	Quality	Salts	PO ₄	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 0.3 mg/l
						Pathogens	E-coli	domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	0 counts / 100 ml
						Pathogens	Total Coliform	domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	<10 counts / 100ml
						Abstraction	Seasonal abstraction: water level recovers from abstraction impact during wet season, under consideration of climate change and drought cycles. Permanent abstraction: water level decline stabilises under consideration of aquifer response time.	Groundwater use should be sustainable for all users and the environment	n/a
				Groundwater (all)	Quantity	Discharge	Relative water levels between groundwater and surface water (in mamsl)	The natural gradient between groundwater and surface water should be maintained	n/a
							Buffer zones	No groundwater abstraction around wetland and river FEPAs in accordance with the implementation manual for FEPAs.	250m
				Groundwater (Cenozoic coastal sand)		Nutrients	NO ₃ (as N)	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 8.2 mg/l
						Salts	EC	domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 520 mS/m
				Groundwater (Basement)	Quality	Nutrients	NO ₃ (as N)	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 11.0 mg/l
						Salts	EC	domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 899 mS/m
				Groundwater (all)		Salts	PO ₄	domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 0.3 mg/l
						System variable	pH	domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	6.7 - 8.3
						Pathogens	E-coli	domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	0 counts / 100 ml
							Total Coliform	domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	<10 counts / 100ml

IUA	Class	Quaternary Catchment	RU	Resource Name	Component	Sub Component	Indicator/ Measure	RQO Narrative	RQO Numeric
A3 West Coast	III	G21B	9-Atlantis	Groundwater (all)	Quantity	Abstraction	Seasonal abstraction: water level recovers from abstraction impact during wet season, under consideration of climate change and drought cycles. Permanent abstraction: water level decline stabilises under consideration of aquifer response time.	Groundwater use should be sustainable for all users and the environment	n/a
						Groundwater level	Water level	Minimum water level in abstraction boreholes within 2.5km from the ocean to avoid saline intrusion	>1 mamsl
						Discharge	Relative water levels between groundwater and surface water (in mamsl)	The natural gradient between groundwater and surface water should be maintained	n/a
							Buffer zones	No groundwater abstraction around wetland and river FEPAs in accordance with the implementation manual for FEPAs.	250m
						Nutrients	NO3 (as N)	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 2.3 mg/l
				Groundwater (Cenozoic coastal sand) Groundwater (Basement) Groundwater (all)	Quality	Salts	EC	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 287 mS/m
						Nutrients	NO3 (as N)	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 10.4 mg/l
						Salts	EC	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 1052 mS/m
						System variable	pH	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	6.7 – 8.3
						Pathogens	E-coli	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	0 counts / 100 ml
D10 Diep	III	G21D	10-Malmesbury	Groundwater (all)	Quantity	Pathogens	Total Coliform	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	<10 counts / 100ml
						Abstraction	Seasonal abstraction: water level recovers from abstraction impact during wet season, under consideration of climate change and drought cycles. Permanent abstraction: water level decline stabilises under consideration of aquifer response time.	Groundwater use should be sustainable for all users and the environment	n/a

IUA	Class	Quaternary Catchment	RU	Resource Name	Component	Sub Component	Indicator/ Measure	RQO Narrative	RQO Numeric
E12 Cape Flats	III	G22C, G22D, G22E	2-Cape Flats	Groundwater (all)	Quantity	Discharge	Buffer zones	No groundwater abstraction around wetland and river FEPAs in accordance with the implementation manual for FEPAs.	250m
						Low flow in river	Compliance with the low flow requirements in the river (as per riverine RQO)	Maintain (groundwater component of) the low flow requirements in the river	Maintenance low flow requirements: 0.578 (6.22 %MAR) at node Biv6 (no gauge)
						Discharge	Relative water levels between groundwater and surface water (in mamsl)	The natural gradient between groundwater and surface water should be maintained	n/a
						Nutrients	NO3 (as N)	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 7.1 mg/l
						Salts	EC		< 358 mS/m
						Nutrients	NO3 (as N)		< 6.4 mg/l
						Salts	EC		< 617 mS/m
						System variable	pH		6.3 – 8.6
						Pathogens	E-coli		0 counts / 100 ml
						Pathogens	Total Coliform		<10 counts / 100ml
						Groundwater level	Water level	Minimum water level in abstraction boreholes within 2.5km from the ocean to avoid saline intrusion	>1 mamsl
						Discharge	Buffer zones	No groundwater abstraction around wetland and river FEPAs in accordance with the implementation manual for FEPAs.	250m
						Low flow in river	Compliance with the lowflow requirements in the river	Maintain (groundwater component of) the low flow requirements in the river, as per surface water RQO requirement	Maintenance low flow: 0.348 Mm3/a (7.74 %MAR) at Bvii7 (no gauge)

IUA	Class	Quaternary Catchment	RU	Resource Name	Component	Sub Component	Indicator/ Measure	RQO Narrative	RQO Numeric
				Superficial aquifers	Quantity	Discharge	Relative water levels between groundwater and surface water (in mamsl)	The natural gradient between groundwater and surface water should be maintained	n/a
				Groundwater (Cenozoic coastal sand)		Nutrients	NO3 (as N)		< 9.2 mg/l
						System variable	pH		6.6 – 8.4
						Salts	EC		< 180 mS/m
				Groundwater (Basement)	Quality	Nutrients	NO3 (as N)	Groundwater should be fit for domestic use after treatment; and groundwater quality shall not show a deteriorating trend from natural background	< 11.0 mg/l
						Salts	EC		< 953 mS/m
				Groundwater (all)			E-coli		0 counts / 100 ml
						Pathogens	Total Coliform		<10 counts / 100ml